



## Pediatric Sample Requirements

Allele Diagnostics  
 120 N. Pine St, Suite 152  
 Spokane, WA 99202  
 Phone: 844-255-3532  
 Fax: 509-232-5779  
 Email: info@allelediagnostics.com

MICROARRAY		
Sample Type	Quantity Required	Comments
Peripheral Blood	2-3 mL NaHep and 2-3 mL EDTA whole blood Neonates: 1 mL NaHep and 1 mL EDTA whole blood	Sufficient for multiple tests Minimum 0.5 mL blood in NaHep
Cord Blood	2-3 mL NaHep and 2-3 mL EDTA whole blood	MCC testing recommended
Buccal Swab	OCD-100 ORAcollect (2 swabs)	Secondary specimen may be needed for FISH visualization
DNA	5 µg extracted in a CLIA-certified lab	Recommend Puregene extraction kit
KARYOTYPE		
Sample Type	Quantity Required	Comments
Peripheral Blood	3-5 mL NaHep whole blood Neonates: 1 mL NaHep whole blood	Minimum 0.5 mL blood
Cord Blood	2-3 mL NaHep whole blood	
SEQUENCING / MOLECULAR		
Sample Type	Quantity Required	Comments
Peripheral Blood	2-3 mL EDTA whole blood Neonates: 1 mL EDTA whole blood	
Cord Blood	1 mL EDTA whole blood	MCC testing recommended
PARENTAL SAMPLES		
Sample Type	Quantity Required	Comments
Peripheral Blood	2-3 mL NaHep and 2-3 mL EDTA whole blood	
Buccal Swab	OCD-100 ORAcollect (2 swabs)	For microarray, MCC, or sequencing <b>only</b>

### Additional Comments:

- Sodium heparin (NaHep) = green top tube; EDTA = lavender top tube
- Heel sticks: Not an acceptable specimen type.
- Lithium heparin: Sodium heparin preferred, but lithium heparin is an acceptable alternative anywhere sodium heparin is listed.
- For sample types not listed above, please contact Allele Diagnostics to discuss.

### Shipping Instructions:

- All tubes must be labeled with patient name and date of birth. Enclose completed Allele Diagnostics Pediatric requisition form(s) and any previous cytogenetic/molecular reports.
- Ship at room temperature in a leak-proof, rigid container with overnight delivery.
- If shipping on Friday, mark Saturday delivery on the shipping label.